



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

14 Poles, or $69\frac{1}{2}$ *English* miles and 14 Poles; 8 Furlongs to a mile, and 40 Poles to a Furlong. Which being compared to that measure of a Degree, which is deliver'd in the above-mention'd *French* Discourse, will be found to come very near it, they finding 73 miles *ferè*, at 5000 feet to an *English* mile, which make 365000 feet; whereas the $69\frac{1}{2}$ *English* miles and 14 Poles, found by Mr. *Normood*, amount to 367200 feet, reckoning 5280 feet to an *English* mile, as the true measure of it is; whence the difference between these two measures appears to be no more than 2200 feet, which is not half an *English* mile by 440 feet.

If any one desire to know further the whole *Circumference*, as also the *Diameter* and *Semidiameter* of the said Terraqueous Globe, according to this measure, he will easily find,

The Circumference to be	25056 <i>ferè</i> .
The Diameter,	7966
The Semidiameter,	3983

Observations made of the late Solar Eclipse on the first of June, 1676. st.v.

One, by *Francis Smethwick* Esquire, as followeth:

I *Nitium defectionis* Westmonasterii h.7. 50'. *post med. noctem*
Finis, h 9. 54 $\frac{3}{4}$. *Junii 1. 1676.*

Totius Eclipsis duratio, hora 2. 4 $\frac{3}{4}$.

Tempus observatum fuit cum horologio oscillatorio, vibrante minuta secunda, & correcto per observationes. Tubus adhibitus fuit bonæ notæ, pedum 7 $\frac{1}{2}$.

The other, by Mr. *Colson* at *Wapping*, near *London*, as followeth;

<i>Temp. juxta</i> <i>horol. oscill.</i>	<i>Phases.</i>	<i>Solis</i> <i>alt.</i>	<i>Tempus correct.</i> <i>ex altit.</i>
h. . . "		0	h. . . "
7.34.50		22.46	7.36. 0
7.37.14		33.10	7.38.40
7.39.10	dig.	33.30	7.40.48
7.50.40	$\frac{1}{4}$	—	7.51.51 <i>Tubo optico æstim.</i>
dub. 8. 8.34	$1\frac{1}{4}$	—	8. 9.45 <i>Tubo optico mensur.</i>
8.17.25	$2\frac{1}{5}$	—	8.18.36
8.27.10	$3\frac{1}{10}$	—	8.28.21
9.39.—	$1\frac{1}{2}$	—	9.40.— <i>Tubo æstim.</i>
9.43.—	$1\frac{1}{6}$	—	9.44.—
9.48.—	$\frac{1}{4}$	—	9.49.—
9.54.25	<i>non finita</i>	—	9.55.36
9.55.55	<i>finita.</i>	—	9.57. 6
4.26. 5	<i>Solis alt.</i>	32.10	4.26.56
4.28.58		31.53	4.29.52
4.31. 21		31.31	4.32.16